

**REPORT DOCUMENTATION PAGE***Form Approved*  
**OMB No. 0704-0188**

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

**1. REPORT DATE (DD-MM-YYYY)**

15-06-2001

**2. REPORT TYPE**

UU

**3. DATES COVERED (From - To)****4. TITLE AND SUBTITLE**

See the abstract below

**5a. CONTRACT NUMBER****5b. GRANT NUMBER****5c. PROGRAM ELEMENT NUMBER****6. AUTHOR(S)**

See the abstract below

**5d. PROJECT NUMBER****5e. TASK NUMBER****5f. WORK UNIT NUMBER****7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)****8. PERFORMING ORGANIZATION REPORT NUMBER****9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)****10. SPONSOR/MONITOR'S ACRONYM(S)****11. SPONSOR/MONITOR'S REPORT NUMBER(S)****12. DISTRIBUTION / AVAILABILITY STATEMENT**

**DISTRIBUTION STATEMENT A**  
Approved for Public Release  
Distribution Unlimited

**20010705 171****13. SUPPLEMENTARY NOTES****14. ABSTRACT**

Phone.txt



DTICPUB0615.txt

US NAVAL OBSERVATORY COLLECTION

**15. SUBJECT TERMS****16. SECURITY CLASSIFICATION OF:**  
UU**17. LIMITATION OF ABSTRACT****18. NUMBER OF PAGES****19a. NAME OF RESPONSIBLE PERSON**  
See the abstract above**a. REPORT****b. ABSTRACT****c. THIS PAGE****19b. TELEPHONE NUMBER** (include area code)  
See the phone list above

Mars," Association of Lunar and Planetary Observers Internet Web  
Page: The Mars Section,  
<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/discrete.htm>

Beish, J. D. (2000). "A German Equatorial Mount for the Planetary  
Telescope," The Digital Lens: Newsletter of the Association of  
Lunar and Planetary Observers Computing Section,  
<http://www.m2c3.com/alpocs/tdl2000/mountmath05222000/mount.htm>

Beish, J. D. (2000). "Martian Volcanoes on HST Images," Association  
of Lunar and Planetary Observers Internet Web Page: The Mars  
Section,  
<http://www.lpl.arizona.edu/~rhill/alpo/marstuff/VOLCANO.htm>

Beish, J. D. (2000). "Measuring Celestial Dimensions with Micrometers,"  
The Digital Lens: Newsletter of the Association of Lunar and  
Planetary Observers Computing Section,  
[http://www.m2c3.com/alpocs/tdl2000/celestialdimensions01232000/  
celestial\\_dimensions.htm](http://www.m2c3.com/alpocs/tdl2000/celestialdimensions01232000/celestial_dimensions.htm)

Beish, J. D. (2000). "Nuts and Bolts of Computing the Ephemeris - Part  
Four," The Digital Lens: Newsletter of the Association of Lunar and  
Planetary Observers Computing Section,  
<http://www.m2c3.com/alpocs/tdl1999/nutsbolts08301999/TDL-4.html>

Beish, J.D. (2000). "Practical Calculations for the Newtonian Secondary  
Mirror," The Digital Lens: Newsletter of the Association of Lunar  
and Planetary Observers Computing Section,  
<http://www.m2c3.com/alpocs/tdl2000/telescopemath06202000/field1.htm>

Beish, J. D. (2000). "Some of the 'Astronomical Seeing' Scales," The  
Digital Lens: Newsletter of the Association of Lunar and Planetary  
Observers Computing Section,  
[http://www.m2c3.com/alpocs/tdl2000/seeing01212000/The%20Seeing%  
20Scales.htm](http://www.m2c3.com/alpocs/tdl2000/seeing01212000/The%20Seeing%20Scales.htm)

Hajian, A. R. (1999). "Stellar Surface Structure," in Michelson  
Interferometry Summer School, held at the California Institute of  
Technology, in Pasadena, California, 9-13 August 1999,  
<http://sim.jpl.nasa.gov/michelson/viewgraphs/HajianFinal.PDF>

Hartkopf, W. I. (1999). "Five Reasons why the Most Interesting, Most  
Exciting, and Most Important Objects to Observe  
(Interferometrically or Otherwise) are Binary Stars," in Michelson

Pkg14

Interferometry Summer School, held at the California Institute of  
Technology, in Pasadena, California, 9-13 August 1999,  
<http://sim.jpl.nasa.gov/michelson/viewgraphs/hartkopf.pdf>

Five (5) Reasons  
why the  
*Most Interesting,*  
*Most Exciting,*  
and  
*Most Important*  
OBJECTS TO OBSERVE  
(Interferometrically or Otherwise)  
are  
**Binary Stars**

## **FIVE REASONS ...**

1. Binaries as Scales
2. Binaries as Yardsticks
3. Binaries and Stellar Evolution
4. Binaries in Other Guises
5. Binaries as "Vermin"

Current status of binary star observations

## Reason 1: Binaries as Scales

- Mass is **THE** fundamental quantity — determines luminosity, size, lifetime, heavy element generation, ultimate fate.
- Need binaries to get masses!

But why is interferometry important in binary star work? A two-part answer:

**Part 1:** No single observing technique yields all necessary information

*Example:* astrometric or “visual” orbit  $\rightarrow P, a'', T, e$ , plus orientation angles  $i, \Omega, \omega$

But Kepler’s Third requires linear separation  $a$

Spectroscopic orbit  $\rightarrow P$  and  $a \sin i$  ( $a_1 \sin i$  and  $a_2 \sin i$  if SB2)

Therefore need complementary techniques.

Distance + astrometric orbit  $\rightarrow a \rightarrow$  mass sum

Particularly useful: spectroscopic + astrometric (yields individual masses if SB2)

**Part 2:** Different observing techniques results in different separation or period regimes


- Astrometry: wide, long-period systems
- Spectroscopy: close, short-period systems

Improvements in spectroscopic techniques (coravel, other cross-correlation techniques) → measure smaller RVs → longer periods


Human lifespan limitations, however! Most improvement must come from “visual” side


- Speckle: tens of mas → periods years to decades (25+ years' data)
- Mark III: periods weeks to years (bright stars, small numbers)
- NPOI: periods days (bright stars, just starting)


But why get masses?












# Masses from Speckle Data



1

## Reason 2: Binaries as Yardsticks

Spectroscopic + astrometric orbits  $\rightarrow a'' + a \rightarrow$   
distance ("orbital parallax")

Technique independent of spectral type, distance  
(sort of); works for stars for which trigonometric  
parallax doesn't

### Reason 3: Binaries and Stellar Evolution

A few questions:

- What role does duplicity play in stellar evolution?
- Are ALL stars created in sets of 2 or more?
- Do all stars have a choice — either companions or planetary systems?  
Can they have both?
- Do stars of all spectral classifications show similar duplicity rate?
- How does duplicity change with time — i.e., once formed, how often are binaries disrupted?

Standard number: ~half of stars binaries

WDS: 450,000+ observations, ~80,000 stars,  
200+ years. Sounds pretty good!

Surveys incomplete, however — true numbers not very well known!

- BSC: new “naked eye” stars found by speckle!  
Still 2/3 unchecked
- Hipparcos: 3,500 new binaries (many are observable visually)
- Surveys of stellar samples, but by no means thorough

Problem even worse — need complementary surveys for different separations.

Result: very few attempted.

## One Tantalizing Survey Result

- PMS stars in young star-forming regions (ex.: Taurus-Aurigae, age 0.002 Gyr) have multiplicity rates  $\sim$ twice that for older ( $\sim$ 5 Gyr) solar-neighborhood counterparts. Hyades (0.7 Gyr) rate in between
- Leonard: binary-binary collisions in clusters and associations might eject stars, decrease their duplicity frequency compared to field stars
- Speckle of O stars: find lower frequency for cluster stars than field stars

Little known for  $0.7 < \text{age} < 5$  Gyr — when do ejections occur?

Mason et al: surveyed  $\sim$ 200 solar-type stars (speckle plus micrometry). Ages from chromospheric activity. Find duplicity fraction for more active stars (age  $\sim$  1 Gyr) about 18%, that for less-active stars ( $\sim$ 4 Gyr) 9%.

Need larger sample, data at smaller and larger separations.



## Reason 4: Binaries in Other Guises

Effects of duplicity not always obvious!

*Example:*  $\lambda$  Boo variables:

- Weak metal lines (esp. Mg II)
- C, N, O, S nearly solar
- Most have moderate to high projected rotational velocities
- Types of stars?

Farraggiana & Bonifacio: find  $1/4 - 1/3$  show duplicity (most from speckle + Hipparcos)

Hypothesize most  $\lambda$  Boo stars actually normal binaries

How many types of variables thought due to duplicity? From Sterkin & Jaschek:

- **Eruptive variables:**

1. RS CVn: close binaries with H and K Ca II in emission
2. IN(YY): matter-accreting Orion variables

- **Eruptive supernovae and cataclysmic variables:**

1. Novae (massive white dwarf/cool dwarf binaries):  
include fast, slow, very slow, recurrent types
2. Nova-like systems (WD+WD, WD+MS, etc): include  
AM CVn, AM Her, DQ Her, UX UMa, VY Scl systems
3. Type I supernovae
4. Dwarf novae or U Gem variables: include SS Cyg, Z Cam,  
SU UMa, and Z And or symbiotic stars

- **Eclipsing variables:**

1. EA: Algol types ( $N = 710 - 2000$ )
2. W Ser systems: long-P Algol-like mass-transferring binaries
3. EB: Beta Lyr types ( $N = 706 - 1500$ )
4. EW: W UMa types ( $N = 88 - 1000$ )
5. GS: have one or more giant components
6. PN: one component is nucleus of PN
7. WD: have white dwarf component
8. WR: have Wolf-Rayet component
9. AR: AR Lac type detached systems
10. DM: detached MS systems
11. DS: detached systems with subgiant
12. DW: detached systems like W UMa system
13. KE: contact systems of early spectral type
14. KW: contact systems of late spectral type
15. SD: semi-detached systems

- **X-ray sources:** 9 categories of bursters, novae, pulsars

What can interferometry contribute?

- Sizes, shapes of components, hot spots, dark spots, limb-darkening, etc. (other speakers)
- Masses + distances → true for other variables in binaries, as well
- Orbital inclination → trajectory during eclipses; aid study of extended atmospheres, accretion disks, etc.
- Orbital precession → study longer-term photometric, spectroscopic changes

## **Reason 5: Binaries as "Vermin"**

Some people despise binary stars!  
(poor misguided fools)


Reasons: need calibration point sources, guide stars for satellites, missiles, etc.

*Example:* SIM: needs 6,000 grid stars stable to  $4 \mu\text{as}$  over 5 years

Advantage of interferometry over spectroscopy for surveys -- one shot!

Current state of affairs - some good, some bad

# Binary Star Measures in the WDS, 1800-2000



Median Separations of WDS Binaries

